U O P A P P
A F A I P

Author: Bryan Clifford Anderson, DNAP, CRNA

L O :

Upon completion of this study, the CRNA will be able to:

1. Discuss the significance of remifentanil use as a primary analgesic agent in parturients for whom neuraxial anesthesia is not an option.

2. Identify clients in which neuraxial anesthesia might be contraindicated.

3. Compare the efficacy of select medications in the control of pain in paturients.

4. Appraise findings in current evidence to determine to what extent

5. Explain dosing and timing parameters that warrant consideration in the administration of remifentanil for the treatment of labor pain.

In order achieve optimal patient outcomes in anesthesia patients, it is important to consider multiple options for pain control, especially when traditional options pose a problem, or are not options. In particular, there are parturient clients for whom the use of neuraxial anesthesia (epidural and spinal blockade) is not an option. In these case an alternative option, that warrants consideration for patient centered anesthesia practice is the use of remifentanil (ultiva). Guidelines for the use of remifentanil in obstetric patients are sparse, poorly developed and are not readily available to anesthesia practitioners.

P A L

There is no question about the amount and extent of pain associated with child birth. There are some common interventions used to ameliorate pain, including the use of epidural anesthesia. However, there are several reasons that an epidural may be contraindicated during labor including the presence of coagulopathies, anticoagulation therapy, prior back surgery, patient refusal, or the inability to safely place an epidural. In labor patients for whom neuraxial anesthesia is not an option, there are limited alternative choices that have been explored or considered. Once such possibility deserving of consideration is the use of the opioid remifentanil as the primary analgesic for the management of pain associated with labor.

Olufolabi et al identified "that the cyclical pattern of labor pain, as compared with continuous postoperative surgical pain, would benefit from bolus delivery of a short-acting drug that produced its analgesic effect only during contractions and was without significant maternal and fetal side effects." [1] One such drug that should be considered is remifentanil (ultiva).

I E P P O P L P

The use of remifentanil in the parturient as the primary analgesic is significant for several reasons; the most salient of which is the basic human right to the management of pain. Pain, as defined by the International Association for the Study of Pain (IASP), is "an unpleasant sensory and emotional experience associated with actual or potential tissue damage."

be addressed and managed in accordance with the needs and wishes of the individual patient. Interventions that alleviate or eliminate pain are not merely a matter of beneficence, but also form part of the duty to prevent harm.[3]

The variations in pain perception among women in labor creates an essential component in the administration of anesthesia in the provision of patient-centered anesthesia care. One recent study identifies that the perception of labor pain was equivalent to a digit amputation without anesthesia.[4] Even though variability regarding the intensity of pain exists among women during labor, the majority of women do experience more than minimal pain during this time.[5] Negative psychological effects of pain associated with labor can occur in some women. "Psychological harm can be experienced through the provision or withholding of labor analgesia, underscoring the tremendous variability in the meaning of labor pain for different women." [5] Interventions to alleviate pain in labor have effects on much more than the physical aspects of pain, but also include the emotional and psychological factors.

Epidural analgesia is considered the standard for pain management during labor.[5] Access to pain management is a right that is fundamental and should not be withheld or denied to any patient regardless of age, ethnicity, or socioeconomic status. This right is violated if a parturient is unable to partake in standard methods used for managing the pain of labor. The significance of analgesia during labor is related to the access parturients have to care. "Equity is concerned with maximizing fairness in the distribution of healthcare services… and minimizing disparities in health."[6] By utilizing an intervention such as remifentanil, the alleviation of pain associated with labor encompasses a greater portion of this population.

A patient's perception of their analgesic regimen is also a concern. It is central for a provider to address this intervention that is rooted in reliable evidence. The use of patient-controlled analgesia (PCA) puts the patient in control when dosing of medication occurs and is considered the gold standard for acute pain management.[7] Patient satisfaction with remifentanil as a primary analgesic for labor pain is an important topic within this context. There is evidence that maternal satisfaction is influenced by factors other than age, ethnicity, socioeconomic status, pain, medical interventions, and continuity of care, when women evaluate their childbirth experiences. These overriding factors have been identified as personal expectations, the amount of support

tightness. With remifentanil use, profound analgesia may be achieved with minimal effect on cognitive function, and low doses of remifentanil can be used to maintain anesthesia in spontaneously breathing patients.[15, 17]

Remifentanil is licensed for induction and maintenance of general anesthesia; however, it is currently an 'off-label' use in obstetrics.[17] Even though remifentanil is not licensed for use in obstetric patients, administration of drugs outside their product license is a common occurrence in obstetric anesthesia.[18]

C P P N P P A P I P N -

Several factors must be considered when discussing the use of neuraxial anesthesia for labor analgesia. These include generally recognized absolute and relative contraindications for neuraxial anesthesia such as bleeding or clotting disorders (coagulopathies), severe hypovolemia, elevated intracranial pressure, valvular heart disease, infection at injection site, or patient refusal.[10] Additional factors include anticoagulation therapy, prior back surgery, or the inability to perform a neuraxial anesthetic. This discussion is not intended to be all-inclusive, but rather to highlight several clinically relevant factors regarding the

including potential for short or long-term complications, fear of pain with implementation, fear of numbness or altered sensation, lack of control over the anesthetic, or the inability to obtain adequate anesthesia. Proper preparation, education, and collaboration are keys to successful interaction with patients.[10] In cases where a parturient declines the use of neuraxial anesthesia techniques, the provider must be prepared to offer an alternative for managing the pain associated with labor. Alternatives provide access to pain management, and uphold the fundamental right that pain management should not be withheld or denied to any patient.[3]

Reflection Box 3. Neuraxial anesthesia techniques vs. alternative anesthesia methods

1. Consider cases in which you used Neuraxial anesthesia techniques.
2. Have there been cases in which would have considered alternatives to neuraxial anesthesia?
3. What were the factors that prompted a desire for an alternative technique?

[illegible]

The literature that compiled and reviewed in the most current studies (meta-analysis, systematic review, reviews of literature, and focused review) revealed that the use of remifentanil as a primary analgesic for the management of labor pain is an accepted practice. When implemented appropriately (regardless of methodology), it is more effective than IV meperidine but less effective than an epidural.[23-26] A 2010 Cochrane Review investigated different parenteral opioids for maternal pain relief during labor and concluded that there is insufficient evidence to identify the best opioid for pain relief.[27] (Insert Table 1 here)

An Overview of Remifentanil (Ultiva)

Classification and Metabolism	Intravenous opioid with rapid onset and brief duration. Regulated as a schedule II substance	The brief duration results from rapid metabolism by plasma and tissue esterases, and not from hepatic metabolism or renal excretion.
Potency	100 times more potent than morphine.	Fentanyl is also 100 times more potent than morphine.
Administration and Duration	Administered via continuous intravenous infusion	Effects begin in minutes and end 5 to 10 minutes once stopped.
Common Dose	*For Surgical Anesthesia*: 0.05 to 2 mcg/kg/min (Current evidence varies)	*For Post-Operative Anesthesia*: 0.025 to 0.2 mcg/kg/min (Current evidence varies)
Adverse Effects	*During Infusion*: respiratory depression, hypotension, bradycardia, and muscle rigidity sufficient to compromise breathing	*Post infusion*: nausea (44%), vomiting (22%), and headache (18%).

Adapted From: Lehner RA. (2013) Pharmacology for nursing care (8th Ed.). Elsevier: St. Louis

In a 2012 meta-analysis by Schnabel et al evaluating the efficacy of remifentanil PCA compared with other techniques for labor analgesia, 12 randomized controlled trials with a total of 593 participants, 269 of which received remifentanil, were included. Of the 12 trials in the meta-analysis, healthy term parturients (ASA classification I and II) without a history of opioid use, drug abuse, allergy to remifentanil or abnormal hepatic or renal function were included. Four different active comparators were investigated– meperidine, fentanyl, nitrous oxide, and epidural analgesia. Due to limited data, the authors were only able to pool data for the comparison between remifentanil and either meperidine or epidural analgesia.[23]

Eight trials compared remifentanil with meperidine in this meta-analysis; 208 parturients received a remifentanil PCA and 209 received meperidine either via PCA, as a continuous infusion, or as an intramuscular injection. In allauhP authors weist

(P<0.001, 95% CI 19 to 31mm). When all seven studies were included in the meta-analysis, remifentanil reduced the mean VAS scores at 1 hour by 25mm compared with meperidine (95% CI 20 to 29mm). Compared with the sensitivity analysis of all 7 studies, the summary estimate of the 3-study analysis was unchanged, although the precision was reduced, as reflected by a wider confidence interval.

The efficacy of meperidine, fentanyl, and remifentanil PCA for labor analgesia varied from mild to moderate in this study. Remifentanil PCA provided better analgesia than meperidine and fentanyl PCA during the first hour of treatment. The authors recommend the use of remifentanil only in the last phase of cervical dilation and with continuous monitoring. Further studies were recommended to determine the safety of remifentanil especially with relation to its respiratory effects.[29]

Another double-blind RCT evaluated was conducted by Evron et al in 2005. Eighty-eight healthy term parturients were enrolled in the study and were randomly assigned to receive either increasing doses of PCA remifentanil or an IV infusion of meperidine. For the 43 parturients randomized to receive remifentanil, each received a bolus of 20mcg as a starting dose, regardless of weight, with a 3-minute lockout interval. The dose was increased every 15 to 20 minutes by 5mcg increments, on patient request, to a maximum dose limit of 1500mcg/hr. The 45 parturients who were randomized to the meperidine group received 75mg of meperidine in 100mL of normal saline over 30 minutes and in case of insufficient analgesia, another dose of 75mg, followed by 50mg when necessary, was administered, to a maximum dose of 200mg of meperidine. [30]

The authors concluded that PCA remifentanil use during labor and delivery was associated with improved VAS scores, higher patient satisfaction, and less need to cross over to epidural analgesia compared to IV meperidine. The use of remifentanil appeared to provide better analgesia than meperidine throughout

fentanyl was administered through the epidural catheter and a continuous infusion of 8ml/hr of 0.125 % levobupivacaine and 2mcg/ml fentanyl was initiated. Further boluses of 5-10ml of 0.125 % levobupivacaine were given upon request. In the CSE group, a needle-through-needle technique was performed with 2mg levobupivacaine and 15mcg fentanyl (total volume of 2mL) injected intrathecally with the epidural catheter inserted and connected to the same continuous infusion used in the epidural group. In the remifentanil group, the PCA device was set to deliver 0.1mcg/kg of remifentanil diluted with saline and given as a solution of 25mcg/mL as a bolus infused during a period of 1 minute, with a lockout time of 1 minute. During the study, the PCA bolus was increased following a dose escalation scheme (0.1–0.2–0.3–0.5–0.7–0.9mcg/kg) after every second contraction until the parturient answered 'no' to the question whether she would like to get more efficient pain relief or until a maximum dose of 0.9mcg/kg was achieved. [32]

No statistically significant differences were observed among the three groups with regard to average VAS score at analgesia request (epidural group 64.5mm ±12.84, remifentanil group 66.4mm ±11.50, CSE group 65.8mm ±12.10, P=0.089). CSE group showed a score of 22.56mm ±7.57 versus 34.3mm ±9.8 for remifentanil and 35.6mm ±10.2 for epidural (P=0.000). The authors concluded in terms of labor duration, average VAS pain scores, and maternal overall satisfaction score with analgesia, CSE analgesia is superior to that provided by epidural analgesia or PCA with remifentanil for pain relief. There were no differences in the mode of delivery, side effects or neonatal outcomes between the three techniques.[32] Other than the method used within the study, no further recommendations regarding remifentanil PCA dosing or implementation were provided.

A randomized clinical trial that compared remifentanil and neuraxial techniques was published by Stocki et al in 2014.[33] The primary objective was to demonstrate noninferiority of remifentanil labor analgesia compared with epidural analgesia in laboring women. Thirty-nine parturients participated with random allocation of 19 in the remifentanil group and 20 in the epidural group. Remifentanil was given as a bolus dose and titrated to effect from 20mcg up to a maximum of 60mcg as required with an initial lockout interval of 2 minutes and no background infusion. The PCA bolus/lockout interval was titrated to an end point of either patient comfort, or a maximal bolus dose of 60mcg/minimal lockout interval of 1 minute. For the epidural group an incremental initial loading dose of 15ml of 0.1% bupivacaine with 50mcg fentanyl was administered followed by patient-controlled epidural analgesia infusion of 0.1% bupivacaine with fentanyl 2mcg/ml. A basal infusion of 5ml/hr, with patient-controlled bolus of 10ml and 20-minute lockout was initiated. Additional epidural bolus doses (either 0.1% bupivacaine 10ml during the first stage of labor or 1% lidocaine 8ml during the second stage of labor) were administered to treat breakthrough pain. [33]

In this study, maternal pain was assessed using an 11-point verbal numerical rating scale (NRS) of 0 to 10, where 0 = no pain and 10 = the worst pain imaginable. There was no significant difference found between baseline NRS pain scores in the two groups. Both remifentanil and epidural analgesia resulted in a significant decrease from baseline NRS scores over time. It was observed that scores were significantly lower at 30 minutes in both groups with change for remifentanil of -4.7 ±0.6 and -7.2 ±0.6 for epidural (P<0.0001). Although both are effective at reducing NRS pain scores, remifentanil is inferior to epidural with regard to the magnitude of the pain score reduction at all time points. Pain

P A

Each of the studies above were analyzed for the remifentanil dosing regimen and implementation method used. The doses, implementation methods, and recommendations found in each study are prga

The overall number of studies specifically investigating the use of remifentanil in parturients are few and most look at only a small fraction of the overall parturient population. Among the specific population of interest- parturients for whom neuraxial anesthesia is not an option- the body of literature contained only a few case studies and these lacked the scientific rigor required to be included in evaluation of this topic. The literature included in this review demonstrated clinical heterogeneity; different study protocols with respect to implementation methods, dosing, timing, rate of administration, lockout intervals, and comparative drugs make it difficult to conduct comparison. Participants in included studies were quite homogeneous in nature with most being healthy ASA 1 or 2 patients who met relatively strict inclusion criteria. This is a very specific body of literature related to the efficacious and safe use of remifentanil as a labor analgesic.

I P P

An issue that was presented in a majority of the studies is that of safety related to remifentanil use in this application. Frequently, assessed parameters that were often a secondary focus of the studies included maternal blood pressure, SPO_2, $ETCO_2$, respiratory rate, and level of sedation. In addition, fetal/neonatal assessment often included Fetal Heart Rate (FHR), umbilical cord pH, and 1 and 5-minute Apgar scores as a measure of assessing adverse response to remifentanil use for labor analgesia. On the maternal side of the safety discussion, most literature suggested that close monitoring of SPO_2, respiratory rate, and level of sedation be undertaken with the use of remifentanil.

In addition to monitoring, the use of supplemental oxygen was also frequently recommended and was implemented in many of the studies. Another frequent recommendation that many authors made was the necessity of having individual nursing care when remifentanil is used in the parturient.[18, 26, 29-31] It is also important to note that many studies investigating the feasibility of remifentanil reported a low number of adverse maternal and fetal events. This low number of adverse events may therefore cause an overestimation of the safety of remifentanil use in labor.[23]

E P C P

The ethical issues surrounding the use of remifentanil for labor analgesia require consideration prior to utilization in clinical practice. The evidence supports the use of remifentanil in the parturient as an acceptable practice. The evidence also supports that the use of this regimen can potentially expand access to pain relief during labor for those who may otherwise be excluded. The relief of pain is a basic human right and as such should not be denied.[3] Remifentanil use meets this need for the relief of pain during labor. The expansion of access and the relief of pain are certainly positive attributes of the use of remifentanil in the parturient.

There are, however, other aspects of ethical concern that each practitioner who accepts the responsibility of providing for a patient requiring or desiring remifentanil as a labor analgesic must take into consideration. It is important to emphasize that every clinical situation in which the use of remifentanil may be utilized requires a thorough evaluation by the practitioner of not only applicable clinical data, but also of the patient and their individual needs during the birthing process.

Not only must the patient be considered in this discussion, but also the unborn child who is wholly dependent upon the physiologic homeostasis provided by the mother. The use of remifentanil as a labor analgesic has proven to be both safe and effective when implemented properly. Several studies have examined the side effects of remifentanil use during the first and second There are, ho

E / -B P

Based upon a thorough review of literature on the subject of remifentanil use in the parturient for whom neuraxial anesthesia is not an option, the use of remifentanil as a labor analgesic is an acceptable practice. The use of remifentanil is considered 'off-label' for obstetric use and there is currently no consensus on the optimal dosing regimen.[17, 18] Large-scale studies with rigorous guidelines and protocols need to be conducted to procure further evidence regarding the optimal use of remifentanil in the parturient. These recommendations would ideally include implementation information and order guidelines for the anesthetist, implementation and usage guidelines for nursing staff, and educational information for patients.

Table 3 displays the recommendations for anesthesia providers that desire to use remifentanil for labor analgesia in the parturient. A PCA bolus of 40mcg with a 2-minute lockout and no background infusion is recommended. Supplemental oxygen should be used in conjunction with continuous monitoring of SPO_2, $ETCO_2$, and cardiotocograph readings. Vital signs (BP, HR, RR, SPO_2, $ETCO_2$, Level of Consciousness (LOC), and Pain) should be documented on a Remifentanil PCA flow sheet every 5 minutes for the first 30 minutes after initiating PCA and then every 15 minutes for the duration of remifentanil use. Individual nursing care should be provided and the nurse should have Advanced Cardiovascular Life Support (ACLS) certification.

Table 3. Putting it all together

Mixture	• Remifentanil 2mg diluted in 50ml Normal saline (40mcg/ml)
Delivery Method	• PCA
Dosing	• 40mcg Bolus • 2 minute lockout
Implementation	• Dedicated IV site for Remifentanil with carrier fluid running at 100ml/hr • O2 via Nasal cannula @ 2-3L/min
Monitoring / Documentation	• Continuous SPO2 monitoring with audible alarm for ≤ 93% • Continuous ETCO2 monitoring with audible alarms for ≥ 55mmHg • Continuous cardiotocograph monitoring • Vitals (BP, HR, RR, SPO2, ETCO2, LOC, Pain) - every 5 minutes for the first 30 minutes after initiating PCA - then every 15 minutes for the duration of remifentanil use • All times and vitals documented on Remifentanil flow sheet • Reconciliation per facility protocol of Remifentanil use and waste
Staffing	• 1:1 nursing care with ACLS trained provider • Supervising Anesthetist in-house and immediately available
Other	• Concomitant use of other opioid analgesics is not recommended

It is further recommended that the Supervising Anesthetist be in-house and immediately available for the entire duration of remifentanil use. Concomitant use of other opioid analgesics is not recommended. These recommendations are an amalgamation of evidence gleaned from the studies analyzed and should not be considered absolute. CRNAs must take into account their own clinical knowledge as well as individual patient needs and desires prior to implementing remifentanil in the parturient.